


KIX 1001: ENGINEERING MATHEMATICS 1

Assignment (10%): Matrix Algebra for Homogeneous & Non-Homogeneous Linear Algebraic System

1. The following linear algebraic equations, $[Q]\{c\} = \{b\}$ represents the 4 dof lake system, where RHS vector consists of the mass loading rate of chloride to each of the 4 lakes and $c_1, c_2, c_3,$ and $c_4 =$ the resulting chloride concentrations for Lakes Powell, Mead, Mohave, and Havasu, respectively.



$$\begin{bmatrix} q_{12} & 0 & 0 & 0 \\ -q_{12} & q_{23} & 0 & 0 \\ 0 & -q_{23} & q_{34} & 0 \\ 0 & 0 & -q_{34} & q_{44} \end{bmatrix} \begin{Bmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \end{Bmatrix} = \begin{Bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{Bmatrix} = \begin{Bmatrix} 750.5 \\ 300 \\ 102 \\ 30 \end{Bmatrix}$$

- If the volumetric flow rates are given as follow: $q_{12} = 13.422, q_{23} = 12.252, q_{34} = 12.377$ and $q_{44} = 11.797$. Use determinant analysis to check the condition of the system, then predict the characteristic of the solution without calculation. (1 mark)
- Solve the concentrations in each of the four lakes by using GEwPP method. The calculation must involve the scaling, partial pivoting, forward elimination and the backward substitution procedures. (2 marks)
- Discuss the advantage of GEwPP method and disadvantage of Cramer's rule in terms of efficiency and accuracy for solving high dimension inverse problem with singular matrix. (1 mark)
- Obtain the eigenvalue matrix and normalized eigenvector matrix of the $[Q]$. Note: Arrange the eigenvalue, λ in ascending order ($\lambda_1 < \lambda_2 < \dots < \lambda_n$). (5 marks)
- By using the result in (d), compute the matrix \mathbf{K} indirectly by simplify the calculation using the following information: $\mathbf{Q} = \mathbf{K}^4, (\mathbf{Q} - \lambda\mathbf{I})\mathbf{c} = \mathbf{0}$, eigenvalue property of $\mathbf{Q}^k = \mathbf{P}\mathbf{D}^k\mathbf{P}^{-1}$. (1 mark)

$$\mathbf{K} = \begin{bmatrix} q_{12} & 0 & 0 & 0 \\ -q_{12} & q_{23} & 0 & 0 \\ 0 & -q_{23} & q_{34} & 0 \\ 0 & 0 & -q_{34} & q_{44} \end{bmatrix}^4 \begin{Bmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \end{Bmatrix}_{\lambda_3}$$

where $\{c\}_{\lambda_3} =$ normalized eigenvector of the 3rd eigenvalue (i. e. 2nd largest)